

## **Effects of Geoduck (*Panopea generosa* Gould, 1850) Aquaculture Gear on Resident and Transient Macrofauna Communities of Puget Sound, Washington**

Author(s): P. Sean McDonald, Aaron W. E. Galloway, Kathleen C. McPeck and Glenn R. Vanblaricom

Source: Journal of Shellfish Research, 34(1):189-202.

Published By: National Shellfisheries Association

DOI: <http://dx.doi.org/10.2983/035.034.0122>

URL: <http://www.bioone.org/doi/full/10.2983/035.034.0122>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

## EFFECTS OF GEODUCK (*PANOPEA GENEROSA* GOULD, 1850) AQUACULTURE GEAR ON RESIDENT AND TRANSIENT MACROFAUNA COMMUNITIES OF PUGET SOUND, WASHINGTON

P. SEAN MCDONALD,<sup>1,2\*</sup> AARON W. E. GALLOWAY,<sup>3</sup> KATHLEEN C. MCPEEK<sup>2</sup> AND GLENN R. VANBLARICOM<sup>2,4</sup>

<sup>1</sup>Program on the Environment, University of Washington, Box 355679, Seattle, WA 98195; <sup>2</sup>School of Aquatic and Fishery Sciences, University of Washington, Box 355020, Seattle, WA 98195; <sup>3</sup>School of the Environment, Washington State University, Albrook 202, Pullman, WA 99164; <sup>4</sup>US Geological Survey, Ecosystems Mission Area, Washington Cooperative Fish and Wildlife Research Unit, School of Aquatic and Fishery Sciences, University of Washington, Box 355020, Seattle, WA 98195

**ABSTRACT** In Washington state, commercial culture of geoducks (*Panopea generosa*) involves large-scale out-planting of juveniles to intertidal habitats, and installation of PVC tubes and netting to exclude predators and increase early survival. Structures associated with this nascent aquaculture method are examined to determine whether they affect patterns of use by resident and transient macrofauna. Results are summarized from regular surveys of aquaculture operations and reference beaches in 2009 to 2011 at three sites during three phases of culture: (1) pregear (–geoducks, –structure), (2) gear present (+geoducks, +structures), and (3) postgear (+geoducks, –structures). Resident macroinvertebrates (infauna and epifauna) were sampled monthly (in most cases) using coring methods at low tide during all three phases. Differences in community composition between culture plots and reference areas were examined with permutational analysis of variance and homogeneity of multivariate dispersion tests. Scuba and shoreline transect surveys were used to examine habitat use by transient fish and macroinvertebrates. Analysis of similarity and complementary nonmetric multidimensional scaling were used to compare differences between species functional groups and habitat type during different aquaculture phases. Results suggest that resident and transient macrofauna respond differently to structures associated with geoduck aquaculture. No consistent differences in the community of resident macrofauna were observed at culture plots or reference areas at the three sites during any year. Conversely, total abundance of transient fish and macroinvertebrates were more than two times greater at culture plots than reference areas when aquaculture structures were in place. Community composition differed (analysis of similarity) between culture and reference plots during the gear-present phase, but did not persist to the next farming stage (postgear). Habitat complexity associated with shellfish aquaculture may attract some structure-associated transient species observed infrequently on reference beaches, and may displace other species that typically occur in areas lacking epibenthic structure. This study provides a first look at the effects of multiple phases of geoduck farming on macrofauna, and has important implications for the management of a rapidly expanding sector of the aquaculture industry.

**KEY WORDS:** aquaculture effects, benthic community, geoduck, habitat provision, macrofauna, press disturbance, structural complexity, geoduck, *Panopea generosa*

### INTRODUCTION

Habitat complexity influences diversity and abundance of species through strong effects on predation (Crowder & Cooper 1982) and competition (Grabowski & Powers 2004), as well as by processes such as recruitment, food delivery, and biodeposition driven by flow and turbulence (e.g., Spencer et al. 1997, Lapointe & Bourget 1999, Lenihan 1999). Placement of structures on soft-sediment substrata is known to initiate a number of physical, geochemical, and ecological processes in the disturbed area (e.g., Wolfson et al. 1979, Davis et al. 1982). Within the conceptual framework of ecological disturbance (*sensu* Pickett & White 1985), placement of structures constitutes a longer lasting or chronic event (i.e., “press” disturbance [Glasby & Underwood 1996]) that may affect a number of ecological functions and processes over long time periods. Organisms that are absent from adjacent unstructured areas may colonize newly available surfaces and interstices, altering species diversity dramatically. Moreover, macroalgae growing on aquaculture structures can further enhance emergent struc-

ture and provide additional biogenic habitat (Powers et al. 2007). These changes may attract mobile consumers, such as transient fish and macroinvertebrates (e.g., Davis et al. 1982), a pattern attributed to enhanced resource supplies for detritivores (e.g., sea cucumbers), herbivores (e.g., urchins and some crab species) and predators (e.g., sea stars and other crab species [Inglis & Gust 2003, Dubois et al. 2007]). Moreover, these structures may serve as refugia that reduce individuals’ predation risk (e.g., Dealeris et al. 2004). Conversely, species that require soft-sediment habitat or prey therein may be excluded when structure additions occur (e.g., Woodin 1981). These disturbances may modify predation pressure and alter patterns of primary production (indirect mediation of top-down control [Genkai-Kato 2007]) and trophic dynamics (Grabowski 2004, Grabowski & Powers 2004).

Projections of future aquaculture production to meet human food demands (Costa-Pierce 2002, Dumbauld et al. 2009) imply an expanding ecological footprint for these activities in near-shore environments. Addition of cultured shellfish (e.g., live animals, shell) and aquaculture gear, including bags, racks, and ropes, may substantially increase structural complexity in soft-sediment habitats where these activities frequently occur, and this can affect resident and transient fish and macroinvertebrates.

\*Corresponding author. E-mail: psean@u.washington.edu  
DOI: 10.2983/035.034.0122

For example, netting used to reduce predation of Manila clams (*Venerupis philippinarum*) in aquaculture operations in the United Kingdom altered patterns of biodeposition, leading to changes in community composition of resident macroinvertebrates, including deposit-feeding polychaetes, consistent with organic enrichment (Spencer et al. 1997). Similarly, Inglis and Gust (2003) observed significantly greater densities of predatory sea stars (*Coscinasterias muricata*) associated with longline mussel farms in New Zealand compared with adjacent reference sites, and scup (*Stenotomus chrysops*) in Narragansett Bay experienced lower disappearance rates (emigration + mortality) at an oyster grow-out site than adjacent areas (Tallman & Forrester 2007). Regardless of the processes involved (e.g., biodeposition or the provision of prey and/or habitat), published literature suggests differences in abundance and diversity at shellfish aquaculture sites relative to unstructured areas (Erbland & Ozbay 2008; see review by Dumbauld et al. [2009]).

Pacific geoducks (*Panopea generosa* Gould 1850; hereinafter geoducks) are the largest burrowing bivalve known (Goodwin & Pease 1987) and range from Baja, California, north to Alaska (Bernard 1983). Aquaculture of geoducks has occurred on a commercial scale since 1996 (Jonathan P. Davis, Taylor Resources Inc., pers. comm. September 13, 2007) and has rapidly developed into an important industry in Washington state and British Columbia, with estimated annual production valued at US\$21.4 million (FAO 2012). Culture practices involve large-scale out-planting of hatchery-reared juvenile clams to intertidal habitats, and installation of PVC tubes and netting to exclude predators and increase early survival. Juvenile clams (shell length, 10–20 mm) are placed in tubes (diameter, 10–15 cm) set vertically in the sediment. Nets typically consist of either small plastic mesh caps stretched over the opening of individual tubes or large, continuous covers over entire plots. Predator exclusion structures are removed after clams reach a size refuge from predators, generally 1–2 y after planting. Clams are harvested after an additional 3–5-y grow-out period (see VanBlaricom et al. [2015] for details).

Although commercial geoduck aquaculture operations boost local economies and increase employment and international trade opportunities, there is a dearth of information regarding potential impacts to nearshore ecosystems. Thus, rapid expansion of geoduck aquaculture operations in intertidal habitats of Puget Sound in Washington state has raised concern among managers, conservation organizations, and the public regarding industry practices that may alter resident ecological communities. In response, the 2007 Washington state legislature passed Second Substitute House Bill 2220, which commissioned a series of scientific studies to “measure and assess” the possible ecological impacts of current practices, including use of predator exclusion structures.

The objectives of the current study were to assess differences in the abundance and diversity of resident and transient macrofauna at sites with (culture) and without (reference) geoduck aquaculture during distinct phases of the aquaculture sequence (prior to gear addition, gear present, and after gear removal). Here, “resident” describes macrofauna species that occupy intertidal beaches throughout their entire benthic life history and demonstrate limited postlarval dispersal, whereas “transient” macrofauna make frequent (often daily, linked to tidal fluctuations in water level) migrations between intertidal and subtidal habitats. The following questions were posed: Do the abundance and diversity of resident and transient macrofauna

differ between culture plots and reference areas? What is the response of the macrofauna community to the addition and subsequent removal of aquaculture gear? The culture plots and reference areas at each site were located close enough to each other (75–150 m) to be considered functionally similar habitats. Evidence of an effect would consist of little or no difference prior to aquaculture, but a distinction between culture plots and reference areas after structures were added. If any differences in resident or transient macrofauna communities were detected when habitat complexity was increased (i.e., when aquaculture gear was present), it was hypothesized that these changes would not persist after gear was removed and the disturbance associated with structure addition was ameliorated.

## MATERIALS AND METHODS

### Study Sites

Work described here was done in South Puget Sound, Washington, a subbasin of Puget Sound composed of those marine waters south and west of Tacoma Narrows (47°16′7.97″ N, 122°33′2.76″ W; Fig. 1 inset). The subbasin is shallow (mean depth, 37 m) and characterized by extensive littoral mud and sandflats (674 km<sup>2</sup>) that constitute more than 15% of the total area (Burns 1985). Because of abundant suitable habitat, South Puget Sound supports substantial commercial culture of bivalves, predominately Pacific oyster (*Crassostrea gigas*), mussel (*Mytilus* spp.), Manila clams (*Venerupis philippinarum*), and most recently geoduck. Three study sites with similar habitat characteristics (Table 1) were selected for this study; Stratford (47°19′10.86″ N, 122°47′38.56″ W) and Rogers (47°14′53.13″ N, 122°49′37.38″ W) are located on the east shore of Case Inlet, and Fisher (47°10′32.28″ N, 122°56′33.79″ W) is located on the south shore of the northeastern portion of Totten Inlet (Fig. 2). None of these sites had been used previously for geoduck aquaculture, which afforded the opportunity to examine the resident and transient macrofauna community prior to the initiation of aquaculture operations (pregear) and the early phases of culture, including the addition of aquaculture structure (gear present) and subsequent removal approximately 2 y later (postgear).

### Surveys of Resident Macroinvertebrates (Infauna and Epifauna)

To investigate the resident benthic macroinvertebrate assemblage at the three study sites, surveys were conducted during low tides (0.5 to –1 m MLLW) from 2009 to 2011 at culture plots and adjacent reference areas. Ten randomly distributed core samples (diameter, 8 cm; depth, 10 cm; surface area, 19.6 cm<sup>2</sup>; volume, 196 cm<sup>3</sup>) were collected in culture plots and adjacent reference areas. In addition, 10 larger excavation samples (diameter, 29 cm; depth, 20 cm; surface area, 660.5 cm<sup>2</sup>; volume, 13.2 L) were taken on each sampling date occurring prior to deployment of protective PVC tubes and nets (pregear), and after removal of the structures (postgear). The small core size was chosen as a cost-effective method for sampling the study plots, and analysis of preliminary samples demonstrated that most benthic infauna were sampled adequately (see VanBlaricom et al. [2015]). Moreover, small cores are used frequently to assess benthic infauna (Simenstad et al. 1991). The excavation samples were used to assess the abundance of larger invertebrates (e.g., sand dollars) that appear infrequently in the smaller cores. Core samples were preserved in 10% buffered formalin solution





Figure 1. Locations of study sites in south Puget Sound, Washington. Inset shows the region of interest; most geoduck aquaculture in Washington state occurs within the area demarcated by the box.

immediately after collection. Excavation samples were sieved (0.5-mm mesh) and enumerated in the field, with retained organisms similarly preserved for laboratory identification when necessary.

Core samples were processed in the laboratory using a standard method of winnowing to extract infaunal organisms (Simenstad et al. 1991, Sobocinski et al. 2010). Freshwater was added to a sample, and the sample was mixed so that

TABLE 1.  
Description of local conditions and biota at geoduck aquaculture sites in Puget Sound (see also Fig. 1).

Site/status	Description	Biota
Stratford site: gear placed June 2009; gear removed April 2011 5,100-m <sup>2</sup> farm, 2,500-m <sup>2</sup> plots	The site is on the east shore of Case Inlet (47°19'10.86" N, 122°47'38.56" W). It has a sandy substrate (grain size, ~500 μm), with a moderate slope from +0.61 m to -0.61 m MLLW. The reference area is 150 m to the south on private property.	Horse clams and cockles are present; sand dollars, patchy.
Rogers site: gear placed November 2008; gear removed April 2011 5,100-m <sup>2</sup> farm, 2,500-m <sup>2</sup> plots	The site is on the east shore of Case Inlet (47°14'53.13" N, 122°49'37.38" W). The substrate is sandy to muddy sand (grain size, ~250–500 μm). The beach is steeper and narrower than other sites. Green algae are abundant, and freshwater seepage occurs. The reference area is 150 m to the south on private property.	Horse clams and cockles are present; graceful crab is abundant; sand dollars, patchy
Fisher site: gear placed June 2009 to July 2009; 90% of gear removed April 2011 2,500-m <sup>2</sup> farm, 2,500-m <sup>2</sup> plots	The site is in the northeast portion of Totten Inlet on the south shore, in the Carlyon Beach area (47°10'32.28" N, 122°56'33.79" W). The substrate is muddy sand (grain size, ~250 μm). The reference area is 75 m to the east on private property.	Horse clams are present; crabs, sea stars, and moon snails are abundant.



**Figure 2.** Summed density of prevalent taxa in scuba surveys of transient macrofauna (fish and invertebrates) defined as species present in at least 10% of surveys. Data were collected on culture plots (Culture) and adjacent reference areas (Reference) at three sites in southern Puget Sound during scuba surveys in 2009 to 2011. Note: The northern kelp crab (*Pugettia producta*) is excluded. Error bars are  $\pm$ SE.

sediments settled to the bottom and the elutriated organisms floated to the surface. Water was decanted through a 500- $\mu$ m sieve and organisms were retained on the collection screen. This process was repeated several times for each sample to ensure all organisms had been separated from sediments. Organisms were identified to species or genus when practical, but in all cases at least to family. Family-level identification has been sufficient to support meaningful quantitative analyses in previous studies (Ferraro & Cole 1990, Dethier 2005). In addition, the processing method just described was used to examine beach spawning by Pacific sand lance (*Ammodytes hexapterus*) opportunistically at study sites during the peak spawning period (November to April). Although our methods did not target spawning specifically (e.g., Moulton & Penttila 2000), winnowing or elutriation has previously been used to assess sand lance spawning because the process of agitating the sample loosens the adhesive eggs from sand grains (Thuringer, unpubl.).

Permutation-based multivariate analysis of variance (PerMANOVA [Anderson 2001]) was used to test for differences in the community data within core samples among plot type (culture plots and reference areas within each site) and phases of culture (pregear, gear present, and postgear) separately for each site (Fisher, Rogers, and Stratford). In addition to the main effects, the interaction of plot type and culture phase was tested, and a significant interaction term was interpreted as evidence that gear addition or removal influenced the community of macroinvertebrate infauna. Thus, evaluation of the interaction term was the principal metric for determining the effect of culture practices. Analyses were conducted in R software (R Development Core Team 2011); significance was set at  $\alpha = 0.05$ .

Distance-based tests for the homogeneity of multivariate dispersion (HMD [Anderson 2006]) were also conducted for further characterization of contrasts of core data between culture plots and reference areas. Homogeneity of multivariate dispersion uses a Bray–Curtis distance matrix of species data to calculate the average distance in multivariate space between individual samples and the calculated centroid of the sample’s

group. The average distance and the associated variability are compared between groups and tested for significance with permutation tests. Caswell and Cohen (1991) hypothesized a positive relationship between multivariate dispersion of samples and disturbance, and previous assessments of disturbance effects have pointed to greater variability of species abundance in samples collected from disturbed areas relative to undisturbed areas when evaluated with HMD (Warwick & Clarke 1993). Because variability is the response of interest in HMD analyses, tests were performed on individual core and excavation samples as the replicated unit; sample averaging would have masked important intersample variability. At each site, HMD analyses were used to test differences between the culture plots and reference areas within each culture phase and within plots across culture phases. Analyses were conducted in R software (R Development Core Team 2011); significance was set at  $\alpha = 0.05$ .

In addition to the community analyses, generalized linear mixed models (GLMMs [McCullagh & Nelder 1989]) were used, assuming Poisson-distributed data, to examine the factors contributing to abundance of selected individual macroinfaunal taxa. In univariate analyses, data from all sites were considered together. The effects of plot type, phase, and their interaction were included, as well as random effects of site and month of sampling. Models were fitted by maximum likelihood assuming a Laplace approximation in the lme4 package (Bates & Maechler 2010) of R software (R Development Core Team 2011). Likelihood ratio tests were used to compare models formally, including the interaction term as part of a “frequentist” hypothesis testing approach. Regression coefficients and their 95% confidence intervals were calculated for each model.

#### Surveys of Transient Fish and Macroinvertebrates

To investigate transient fish and macroinvertebrate assemblages at the three study sites, scuba surveys were conducted during daytime high tides (3–4.25 m above MLLW) from 2009 to 2011. A pair of divers used a metric underwater transect tool adapted from Bradbury et al. (2000) to conduct line transects at each site; each diver surveyed a 1-m swath. Sites were comprised of two 2,500-m<sup>2</sup> habitat spaces: a culture plot with active geoduck farming and a nearby reference area (the same reference area used in the core sampling) with no aquaculture activity. Two 45-m transects were done on each habitat, although there was some variation in transect length, depending on weather conditions and dimensions of the culture plots. Successful surveys were dependent on sufficient water clarity for underwater visibility, coinciding to horizontal Secchi disk measurements of at least 2.5 m. Scuba surveys were conducted monthly from March through August, and bimonthly from September through February.

All observed fish and macroinvertebrates larger than 60 mm were identified and enumerated to species or genus, and observations of size (estimated total length for fish, and diameter, carapace width or length for sea stars, crabs, and other benthic invertebrates), water column position, behavior, and associated substrate type (sand, gravel, tubes + netting, tubes – netting) were recorded. Observed species were assembled into 10 functional groups: sea stars, moon snails, hermit crabs, crabs (Brachyura), other benthic invertebrates, flatfishes, sculpins, other demersal fishes, other nearshore fishes, and sea perch (Table 2). Numbers

TABLE 2.

Functional groups for commonly observed taxa in scuba surveys of three geoduck aquaculture sites in Puget Sound, Washington, 2009 to 2011.

Functional group	Common name	Scientific name	Frequency in surveys (%)
Cockle	Heart cockle	<i>Clinocardium nuttallii</i>	29.6
Crab (true crab)	Graceful crab	<i>Cancer (Metacarcinus) gracilis</i>	89.4
	Kelp crab	<i>Pugettia product</i>	47.0
	Red rock crab	<i>Cancer productus</i>	29.6
	Graceful decorator crab	<i>Oregonia gracilis</i>	7.6
Hermit crab	Black-eyed hermit crab	<i>Pagurus armatus</i>	65.2
	Bering hermit crab	<i>Pagurus beringanus</i>	15.9
Moon snail	Pacific moon snail	<i>Lunatia lewisii</i>	55.3
Other benthic invertebrate	Dendronotid nudibranch	<i>Dendronotus</i> spp.	10.6
	Black-tailed crangon	<i>Crangon nigricauda</i>	4.6
	Giant sea cucumber	<i>Parastichopus californicus</i>	0.8
Sea star	Sunflower star	<i>Pycnopodia helianthoides</i>	53.0
	Pink sea star	<i>Pisaster brevispinus</i>	38.6
	Mottled sea star	<i>Evasterias troschelli</i>	22.7
	Ochre sea star	<i>Pisaster ochraceus</i>	15.9
Flatfish	Speckled sanddab	<i>Citharichthys stigmaeus</i>	42.4
	Starry flounder	<i>Platichthys stellatus</i>	18.9
	Sand sole	<i>Psettichthys melanostictus</i>	6.8
Gunnel	Saddleback gunnel	<i>Pholis ornata</i>	6.1
	Pinpoint gunnel	<i>Apodichthys flavidus</i>	1.5
	Crescent gunnel	<i>Pholis laeta</i>	0.8
Other demersal fish	Plainfin midshipman	<i>Porichthys notatus</i>	4.6
	Sturgeon poacher	<i>Podothecus accipenserinus</i>	5.3
Other nearshore fish	Bay pipefish	<i>Syngnathus leptorhynchus</i>	18.9
	Snake prickleback	<i>Lumpenus sagitta</i>	8.3
	Tubesnout	<i>Aulorhynchus flavidus</i>	0.8
Sculpin	Staghorn sculpin	<i>Leptocottus armatus</i>	37.1
	Roughback sculpin	<i>Chitonotus pugetensis</i>	3.0
Sea perch	Shiner surf perch	<i>Cymatogaster aggregate</i>	6.1
	Striped surf perch	<i>Embiotoca lateralis</i>	0.8

of organisms were converted to raw density values to offset the different transect lengths. Species that occurred in less than 5% of surveys were not included in the data analysis.

Based on observations during SCUBA surveys, it was apparent that many of the transient fish and macroinvertebrates do not occupy intertidal habitats during the winter months (Fig. 2). To reduce the effect of seasonal variability on the abundance of many functional groups, data analysis focused only on the April to September period. Three phases of the aquaculture cycle were represented in the data set: pregear (in 2009, prior to any aquaculture operations [-geoducks, -structure]), gear present (in 2010, during active geoduck aquaculture operation, aquaculture gear in place at culture plots [+geoducks, +structure]), and postgear (in 2011, protective tubes and nets were removed but geoducks remained during grow-out [+geoducks, -gear]). Although the 2010 to 2011 data represent periods in which aquaculture was active, farming occurred at culture plots only; thus, there was no change in epibenthic structure at reference areas.

Data from the three survey sites were not analyzed individually because all sites were considered to have functionally similar habitat for mobile macrofauna. In addition, in some cases the sample sizes would have been smaller than practical for the methods applied if the data were separated by site. Data were (log  $x + 1$ )-transformed in R software with the vegan

package (R Development Core Team 2011), with  $\alpha = 0.05$  for statistical tests of significance.

Analyses of similarity (ANOSIMs [Clarke 1993]) were conducted to assess differences in functional groups between culture plots and reference areas across aquaculture phases. A Bray-Curtis dissimilarity matrix (Bray & Curtis 1957) was used in ranking pairwise combinations of the absolute densities for all functional groups and survey events. Test statistics ( $R$ ) and  $P$  values were generated using Monte Carlo permutation tests with 999 iterations. Values of the  $R$  statistic ranged from -1 to 1, with negative values suggesting larger differences within groups (Clarke & Gorley 2001) and positive values indicating larger differences among groups (McCune et al. 2002). An  $R$  value of zero indicates no differences (McCune et al. 2002).

Visual representations of species abundance in different habitat types and during aquaculture phases were explored using nonmetric multidimensional scaling (NMDS [Kruskal & Wish 1978]). Because NMDS has no assumptions of linearity, it is suitable for any dissimilarity matrix (McGarigal et al. 2000), which makes the procedure useful for visualizing relationships in nonnormal data sets of species abundance (McCune et al. 2002). Nonmetric multidimensional scaling was conducted on a Bray-Curtis dissimilarity matrix of the untransformed, raw density data, and 1,000 iterations were performed to ensure convergence with minimal stress. Stress significance was tested

using a Monte Carlo randomization approach. Linear correlation of the functional groups and NMDS axis scores were used to calculate variable weights. Significant functional groups were determined with permutation tests and were overlaid as vectors on the NMDS plots, which facilitated interpretation of the position of each survey event in ordination space.

Addition of aquaculture gear is a press disturbance (see review by Dumbauld et al. [2009]), and disturbance is generally considered one of the main factors influencing variations in species diversity (e.g., Connell [1978], but see Mackey and Currie [2001]). The Shannon index was used to compare differences in diversity between plots for each aquaculture phase. This measure is commonly used in ecological studies; it combines aspects of species richness and relative abundance to produce a value typically from 0–3.5 (Shannon 1948, Shannon & Weaver 1949). A higher index value indicates greater diversity. Two-sample Welch's *t*-tests (Zar 2010) were used to assess differences in diversity between plots at each stage of geoduck farming.


#### Supplementary Observations of Salmon Smolts

In addition to the fish sampling described earlier, observations were made of salmon smolts in the vicinity of aquaculture operations. Pilot observations by divers and snorkelers indicated that smolts at the study sites were not sampled effectively by those methods, possibly because observers altered fish behavior. Moreover, salmon smolts—in particular, chum (*Oncorhynchus keta*)—typically move along shorelines in shallow water (<2 m [Healey 1979, Simenstad et al. 1982]). Shore-based surveys have been developed as a method of monitoring fine-scale use of shallow nearshore areas by juvenile salmonids (e.g., Young 2009). Concurrent with scuba surveys, shore-based visual surveys were conducted monthly during the spring and summer (March to July) to coincide with out-migration of chum, Pink (*Oncorhynchus gorbuscha*), and coho (*Oncorhynchus kisutch*) salmon smolts (Simenstad et al. 1982). An observer at the water's edge slowly walked along a 50-m transect line parallel to shore, spending 1 min in each 10-m section. Observations were made of all fish encountered up to 5 m offshore. Polarized sunglasses were used when necessary to improve observations. Salmonids were identified to species when possible and enumerated. Additional observations of fish length (total length) and behavior were recorded. On each sampling date, one survey each was completed adjacent to the culture plot and reference area. Successful surveys were dependent on surface conditions, coinciding with a Beaufort scale score of 0–1 (calm or light air).

## RESULTS

#### Surveys of Resident Macroinvertebrates (Infauna and Epifauna)

At all three sites, the community of resident macrofauna consisted primarily of polychaete worms (Annelida), small crustaceans (Arthropoda), and small bivalves (Mollusca). In some locations, echinoids (Echinodermata), larger bivalves, burrowing sea anemones (Cnidaria), and sea cucumbers (Echinodermata) were important community components. All sites were characterized by substantial seasonal variation, and the greatest densities typically occurred during July to September (Fig. 3). Total taxa density in core samples showed substantial site-specific variation, with no consistent pattern of



**Figure 3.** (A–C) Density of total taxa in surveys of resident macrofauna (infauna and epifauna). Data were collected on culture plots (Culture) and adjacent reference areas (Reference) at three sites in southern Puget Sound: Fisher (A), Rogers (B), and Stratford (C). Shaded areas illustrate the aquaculture phase when PVC tubes and nets were in place to protect juvenile geoducks (gear present). Error bars are  $\pm$ SE.

greater density in either culture plots or reference areas across months or sites (Fig. 3). Similar taxa were recorded in cores and excavation samples in most cases. In October 2010, adult sand lance were captured in excavation samples collected at the culture plot and reference area at the Rogers site; densities were  $24.2 \pm 11.9/\text{m}^2$  and  $278.6 \pm 115.7/\text{m}^2$ , respectively. However, subsequent evaluation of core samples revealed no evidence of spawning. No adult sand lance, other forage fish, or fish eggs of any type were observed at the other sites.

In total, 68 taxa from 63 sampling events were collected and identified. Results of the PerMANOVAs illustrate differences in community structure across months of sampling, plot types, and phases at each site (Table 3); however, there were no community-level effects of aquaculture operations as indicated by nonsignificant plot type  $\times$  phase interaction terms (Fisher site: pseudo- $F = 0.049$ ,  $P = 0.116$ ; Rogers site: pseudo- $F = 0.023$ ,  $P = 0.643$ ; Stratford site: pseudo- $F = 0.029$ ,  $P = 0.529$ ).

TABLE 3.

Permutational analysis of variance results for multivariate abundance data for all resident macroinfaunal taxa in core samples.

Site	Factor	df	SS	MS	R <sup>2</sup>	F value	P value
Fisher	Month*	9*	1.269*	0.141*	0.266*	2.2528	0.001*
	Plot*	1*	0.496*	0.496*	0.253*	7.927*	0.001*
	Phase*	2*	0.301*	0.151*	0.047*	2.406*	0.008*
	Plot:Phase	2	0.195	0.098	0.023	1.558	0.116
	Error	27	1.691	0.063	0.411		
	Total	41	3.952				
Rogers	Month*	9*	1.335*	0.1488	0.266*	2.229*	0.001*
	Plot*	1*	1.269*	1.269*	0.253*	19.077*	0.001*
	Phase*	2*	0.236*	0.118*	0.047*	1.770*	0.039*
	Plot:Phase	2	0.113	0.057	0.023	0.848	0.643
	Error	31	2.063	0.067	0.411		
	Total	45	5.016				
Stratford	Month*	9*	2.278*	0.253*	0.398*	2.757*	0.001*
	Plot*	1*	0.792*	0.792*	0.138*	8.623*	0.001*
	Phase*	2*	0.380*	0.190*	0.066*	2.072*	0.020*
	Plot:Phase	2	0.168	0.084	0.029	0.916	0.529
	Error	23	2.111	0.092	0.369		
	Total	37	5.729				

Models included month of sampling (Month), plot type (culture plot or reference area; Plot), phase of culture (pregear, gear present, postgear; Phase), and the interaction of plot type and phase. \* Significant results. Significance was set at  $\alpha = 0.05$ .

Within each site, HMD values for the community data from the pregear phase were similar at culture and reference plots (Table 4). Similarly, there were no significant differences in HMD values for culture and reference plots at any site when aquaculture structures were in place (gear present), although the values were somewhat greater at the Rogers and Fisher sites (Table 4). During the postgear phase, values for culture plots and reference areas were less (relative to the previous phase) and not significantly different at Rogers and Fisher ( $P = 0.335$  and  $P = 0.436$ , respectively). At Stratford, the postgear HMD values for the benthic community were similar to values when aquaculture gear was in place (gear present); however, there was a significant difference in values between the culture plot and reference area ( $P = 0.003$ ; Table 4).

Twelve taxa were selected for univariate analyses using GLMMs based on their frequency in samples (>90%) and presumed ecological importance. Abundance of individual taxa showed marked differences across months, plot type, phases, and the interaction of plot type and phase. Taxa showed no consistent response to geoduck aquaculture. Regression parameter estimates and 95% confidence intervals for GLMMs are included in Figure 4. The abundances of six taxa were affected negatively by geoducks and aquaculture gear, as indicated by a significant plot type  $\times$  phase interaction (GLMM chi square,  $P < 0.05$ ) and negative parameter estimates for the gear-present phase (Fig. 4). However, only two taxa experienced persistent negative effects: the polychaete Families Spiro-nidae (chi square = 22.89,  $df = 2$ ,  $P < 0.001$ ) and Orbiniidae (chi square = 109.17,  $df = 2$ ,  $P < 0.001$ ). Abundance of the amphipod *Americorhium salmonis* (chi square = 174.23,  $df = 2$ ,  $P < 0.001$ ) and polychaete Family Hesionidae (chi square = 341.18,  $df = 2$ ,  $P < 0.001$ ) were reduced by the presence of aquaculture gear but

TABLE 4.

Results of the test of multivariate homogeneity comparing multivariate dispersion (HMD test) of resident macroinvertebrate communities of culture plots and reference areas.

Site	Phase	Multivariate dispersion		F value	P value
		Culture	Reference		
Stratford	Pregear	0.34	0.33	0.007	0.93
	Gear present	0.32	0.35	0.178	0.68
	Postgear	0.35	0.25	14.608*	<0.01*
Rogers	Pregear	0.18	0.19	0.162	0.70
	Gear present	0.28	0.31	0.480	0.69
	Postgear	0.21	0.23	1.026	0.34
Fisher	Pregear	0.20	0.22	0.355	0.57
	Gear present	0.27	0.28	0.261	0.64
	Postgear	0.25	0.22	0.790	0.44

Multivariate dispersion, a measure of  $\beta$  diversity, is associated with environmental stress and disturbance. The measure is calculated as the mean distance of all culture phase/habitat community samples to their group centroid in principal coordinate space as defined by Bray-Curtis compositional dissimilarity. \* Significant results. Significance was set at  $\alpha = 0.05$ .


recovered after gear was removed, and the cumacean *Cumella vulgaris* (chi square = 199.16,  $df = 2$ ,  $P < 0.001$ ) and polychaete Families Glyceridae (chi square = 94.75,  $df = 2$ ,  $P < 0.001$ ) and Opheliidae (chi square = 105.31,  $df = 2$ ,  $P < 0.001$ ) increased during the postgear phase in culture plots relative to reference areas. In addition, the abundance of the polychaete Family Goniadidae (chi square = 10.94,  $df = 2$ ,  $P = 0.004$ ) and anemone Family Edwardsiidae (chi square = 20.505,  $df = 2$ ,  $P < 0.001$ ) increased when gear was present, and recovered to pregear levels after gear was removed. The bivalve genus *Rochefortia* (chi square = 6.99,  $df = 2$ ,  $P = 0.030$ ), nemertean genus *Micrura* (chi square = 0.52,  $df = 2$ ,  $P = 0.772$ ), and polychaete Family Capitellidae (chi square = 4.83,  $df = 2$ ,  $P = 0.089$ ) showed no response to geoduck aquaculture activities.

*Surveys of Transient Fish and Macroinvertebrates*

The presence of aquaculture gear affects the composition of transient fish and macroinvertebrate communities (Fig. 5). No significant differences between culture plots and reference areas were detected by ANOSIM when PVC tubes and nets were absent, either pregear or postgear (Table 5). However, a significant difference was detected between culture plots and reference areas when aquaculture gear was present ( $R = 0.081$ ,  $P = 0.035$ ). Tests of ANOSIM between aquaculture phases (Table 5) resulted in a statistically significant difference when comparing the pregear versus gear-present phases and gear-present versus postgear phases for culture plots ( $R = 0.156$ ,  $P = 0.040$ ;  $R = 0.164$ ,  $P = 0.003$ , respectively). There was also a significant difference between gear-present and postgear reference plots ( $R = 0.090$ ,  $P = 0.029$ ). Low  $R$  values of these tests indicate minimal separation in contrasts between the habitats.

Several two-dimensional NMDS plots were used to aid in visualization of differences between habitats within sites and across phases of aquaculture operations. The NMDS plots also






**Figure 4.** Parameter estimates and 95% confidence intervals for generalized linear mixed models of selected macrofauna. The models included main effects of month of sampling, plot type (geoduck culture or reference area), phase (pregear, gear present, postgear), and their interaction, as well as random effects of site (Fisher, Rogers, and Stratford). As noted in the text, a significant interaction term provides evidence of an effect of aquaculture operations on abundance.

confirmed the assumption that the three sites were functionally similar for purposes of analyzing transient macrofauna communities during April to September. The NMDS ordination of the reference plot data shows some intermixing of sites and clustering of the three sites in multivariate space (Fig. 6). Information on stress, Monte Carlo randomization, and goodness-of-fit testing is included in the caption for each plot (Figs. 6–9).

During 2010, when nets and tubes were used in aquaculture operations (gear-present phase), surveys of culture plots and reference areas were generally separated in ordination space (Fig. 7). Neither habitat type was associated consistently with unique functional groups. However, differences in assemblages between culture plots and reference areas were illustrated by significant vector loadings associated with flatfish, hermit crab, sculpin, sea star, snail, and true crab (Brachyura). True crab

showed weak associations with reference areas overall, whereas sculpin and flatfish correlated highly and were more often associated with reference areas. Two additional NMDS ordination plots represent comparisons of the pregear and gear-present phases (Fig. 8), and the gear-present and postgear phases (Fig. 9).

Survey data for the culture plots when PVC tubes and nets were present were more widely dispersed in ordination space compared with the pregear phase (Fig. 8). Differences in assemblages between pregear and gear-present phases were illustrated by significant vector loadings associated with flatfish, hermit crab, sculpin, sea star, and true crab (Brachyura). Prior to gear deployment, culture plots and reference areas were characterized by flatfish and sea star. Conversely, although communities associated with culture plots were represented by a variety of functional groups when nets and tubes were in place



**Figure 5.** Relative abundance of 10 functional groups of transient fish and macroinvertebrates on geoduck culture plots (Culture) and adjacent reference beaches (Reference) during scuba surveys at three sites in southern Puget Sound (2009 to 2011). Data are presented in three April to October periods comprising three phases: (1) pregear, prior to placement of geoducks or aquaculture gear; (2) gear present, when tubes and nets are in place; and (3) postgear, after nets and tubes have been removed and geoducks are in place.

**TABLE 5.**  
**Results of two-way, crossed analysis of similarity (ANOSIM) tests comparing the transient fish and macroinvertebrate community assemblage in geoduck culture plots and reference areas across three phases of aquaculture operations: pregear, gear present, and postgear.**

Test groups	ANOSIM R	P value
Pregear reference area vs. culture plot	-0.0501	0.761
Gear-present reference area vs. culture plot	0.0808*	0.035*
Postgear reference area vs. culture plot	-0.0254	0.789
Pregear vs. gear-present reference area	0.1176	0.093
Pregear vs. gear-present culture plot	0.1557*	0.040*
Pregear vs. postgear reference area	-0.0268	0.600
Pregear vs. postgear culture plot	-0.0851	0.842
Gear present vs. postgear reference area	0.0900*	0.029*
Gear present vs. postgear culture plot	0.1604*	0.003*

A Monte Carlo permutation test with 999 iterations generated the test statistics (R). \* Significant results. Significance was set at  $\alpha = 0.05$ .

(gear present), flatfish were conspicuously underrepresented. At the same time, reference areas were characterized by flatfish and hermit crab, and less so by true crab and sea star.

In comparisons of gear-present and postgear phases, data from culture plots appear mostly separated in multivariate space, but reference area data overlap and appear more homogenous (Fig. 9). Differences in assemblages between gear-present and postgear phases were illustrated by significant vector loadings associated with clam, flatfish, hermit crab, other nearshore fish, sculpin, and true crab (Brachyura). Of the significant functional groups in Figure 9, true crab and other nearshore fish show the strongest associations with culture plots during the gear-present phase, when PVC tubes and nets were in place.

Species diversity, as calculated by the Shannon diversity index ( $H'$ ), was unaffected by geoduck aquaculture operations

(Table 5). There was no significant difference in diversity between culture plots and reference areas during the phases of culture examined in this study: prior to gear deployment ( $t = 0.703$ ,  $df = 11$ ,  $P = 0.496$ ), gear present ( $t = 0.727$ ,  $df = 18$ ,  $P = 0.476$ ), or after gear had been removed ( $t = 0.309$ ,  $df = 25$ ,  $P = 0.760$ ) (Table 6). Total numbers of organisms observed at culture and reference plots were similar prior to gear deployment (pregear, 2009) and after gear removal (postgear, 2011). However, there was an overall increase in total abundance while aquaculture gear was present, and macrofauna counts were more than two times greater at culture plots compared with the reference areas (Table 5).

**Supplementary Observations of Salmon Smolts**

Salmon smolts, chum (*Oncorhynchus keta*) and Pink (*Oncorhynchus gorbuscha*) salmon, were rarely observed during shore-based visual surveys (total, 8%). When present, schools of salmon traveled parallel to the shoreline in less than 2 m of water. No difference in the occurrence of salmon smolts adjacent to culture plots and reference areas was observed, although evidence is anecdotal, given the low encounter rate. No discernible differences in behavior were observed.

**DISCUSSION**


Resident and transient macrofauna communities respond differently to changes in habitat complexity associated with geoduck aquaculture operations. Although results of the current study suggest that structures associated with geoduck aquaculture have little influence on community composition of resident benthic macroinvertebrates (i.e., nonsignificant plot type  $\times$  phase interaction in PerMANOVA), overall densities of resident epifauna and infauna tended to be lower on culture plots relative to reference areas at two of the three study sites. Resident invertebrate communities were characterized by strong seasonal patterns of abundance and site-specific differences in composition. Dispersion in sample variation, which is commonly used to detect effects of disturbance, did not differ between culture plots and reference areas when aquaculture gear was in place. Some individual taxa responded negatively to the presence of geoducks and aquaculture gear (e.g., polychaete Families Spionidae and Orbiniidae), whereas others responded positively (e.g., polychaete Family Goniadidae and anemone Family Edwardsiidae), and still others were unaffected (e.g., bivalve genus *Rochefortia* and polychaete Family Capitellidae).

**TABLE 6.**

**Results of Shannon diversity index ( $H'$ ) calculations for transient fish and macroinvertebrates at geoduck culture plots and reference areas across three phases of aquaculture operations: pregear, gear present, and postgear.**


Phase	Plot type	Shannon diversity index ( $H'$ )	t-Test results for diversity values	Total organisms observed (n)
Pregear	Reference	1.111	$t = 0.703$ , $df = 11$ , $P = 0.496$	530
	Culture	1.188		628
Gear present	Reference	0.923	$t = 0.727$ , $df = 18$ , $P = 0.476$	795
	Culture	1.021		1,692
Postgear	Reference	1.163	$t = 0.309$ , $df = 25$ , $P = 0.760$	621
	Culture	1.207		694

Differences among culture plots and reference areas were examined with Welch's t-test with  $\alpha = 0.05$ . Total abundance of all observed organisms is included.



**Figure 6.** Two-dimensional nonmetric multidimensional scaling (NMDS) ordination of scuba surveys at reference areas during 2010, which corresponds to when aquaculture gear was in place (gear present) at the culture sites. Stress = 17.24. Stress tested statistically significant with the Monte Carlo randomization approach ( $P < 0.01$ ). A goodness-of-fit Shepard plot showed good correlation between the ordination distances and the Bray–Curtis dissimilarities (linear fit  $R^2 = 0.882$ ).

The paucity of strong effects on the resident macrofauna community (epifauna and infauna) may not be unexpected. Previous studies have suggested that aquaculture effects on benthic infauna are most pronounced in soft-sediment habitats directly below or immediately adjacent to shellfish aquaculture operations as a function of organic enrichment via biodeposition (see the review by Dumbauld et al. [2009]). Interestingly, the two taxa experiencing persistent negative effects of geoduck aquaculture activities—Families Spionidae and Orbinidae—are selective detritivores and deposit feeders, respectively (see Table 1 of VanBlaricom et al. [2015]). In off-bottom aquaculture (e.g., suspended culture), the balance of biodeposition and water flow, which removes deposits, tend to be the strongest determinants of community structure (Mattsson & Linden 1983). In on-bottom aquaculture operations, effects of structural complexity and space competition are difficult to separate from changes in biodeposition (Dumbauld et al. 2009). Quintino et al. (2012) specifically investigated the relative contribution of biodeposition and aquaculture gear (i.e., oyster trestles) and found that structures alone had no effect, whereas biodeposition from sedimentation and organic waste did alter the benthic community. However, Spencer et al. (1997) found that the netting used to reduce Manila clam predation reduced flow and led to changes in benthic community composition consistent with organic enrichment. In the current study, several infaunal taxa recovered to pre-gear abundance, or increased in abundance, after aquaculture gear was removed. Effects on resident macrofauna, particularly infauna and epifauna, may be site specific and likely driven by inherent levels of natural disturbance (Simenstad & Fresh 1995) or flushing (Dumbauld et al. 2009), which may be mediated by aquaculture gear. Physical and chemical variables (e.g., sediment grain size, pore water nutrients) that may contribute to site-specific differences




**Figure 7.** Two-dimensional nonmetric multidimensional scaling (NMDS) plot of scuba surveys at culture plots (solid circles) and reference areas (solid triangles) when aquaculture gear was in place (gear present). The functional group vectors shown are those with  $P < 0.05$ . Stress = 13.87. Stress tested statistically significant with the Monte Carlo randomization approach ( $P = 0.02$ ). A Shepard plot showed good correlation between the ordination distances and the Bray–Curtis dissimilarities (linear fit  $R^2 = 0.925$ ). Vector loadings are shown for significant functional groups ( $P < 0.05$ ).

were not examined in the current study. Thus, elucidating potential mechanisms responsible for differences in the response of infauna requires further study. Additional data and analytical inference would also permit more direct comparison with previous studies done by Spencer et al. (1997), Quintino et al. (2012), and others.

Unlike resident macrofauna, the transient fish and macroinvertebrate community was clearly affected by aquaculture activities. The presence of PVC tubes and nets altered abundance and composition significantly, but not diversity, of transient macrofauna. More than two times more organisms were observed during surveys at the culture plots than at reference areas during the structured phase of geoduck aquaculture, indicating that geoduck aquaculture gear created favorable habitat for some types of Puget Sound macrofauna. Analysis of similarity results demonstrated a statistically significant difference between the transient macrofaunal communities in culture plots and reference areas when aquaculture gear was present (Table 5;  $R = 0.081$ ,  $P = 0.035$ ). Yet, the low  $R$  value of the test suggests minimal ecological difference between the habitats. The NMDS plots provide insight into functional groups that may show preference for culture plots (structured habitat) or reference areas (unstructured habitat) when aquaculture gear is present. In general, true crabs, sea stars, and sea perch were more associated with culture plots, and flatfishes and snails were often associated with reference areas.

The large increase in total abundance of transient macrofauna when aquaculture gear was present suggests that increased complexity afforded by PVC tubes and nets attracted some fish and macroinvertebrates to the habitat. Aggregation of macrofauna to structured habitat, and aquaculture gear in particular, has been well documented (Dealteris et al. 2004, Dubois et al. 2007, Dumbauld et al. 2009). The data from the




**Figure 8.** Two-dimensional nonmetric multidimensional scaling (NMDS) plot of scuba surveys at culture plots and reference areas prior to deployment of aquaculture gear (pregear) and when aquaculture gear was in place (gear present). The functional group vectors shown are those with  $P < 0.05$ . Stress = 14.498. Stress tested statistically significant with the Monte Carlo randomization approach ( $P < 0.01$ ). A goodness-of-fit Shepard plot showed good correlation between the ordination distances and the Bray–Curtis dissimilarities (linear fit  $R^2 = 0.918$ ). Vector loadings are shown for significant functional groups ( $P < 0.05$ ).

current study suggest that provision of foraging and refuge habitat is the primary mechanism for the attraction; crabs and sea stars were frequently observed feeding within culture plots, and smaller fish and crabs were observed retreating under netting when larger animals or divers approached. Similarly, Inglis and Gust (2003) observed increased predation by sea stars at New Zealand longline mussel farms, and Tallman and Forrester (2007) identified refuge value as a major factor leading to greater site fidelity of juvenile scup (*Stenotomus chrysops*) to aquaculture structures in Rhode Island. Increased foraging pressure by transient macrofauna may also provide an additional mechanism to explain slightly depressed densities of resident macrofauna in culture plots relative to reference areas.

In the current study, some taxa, particularly flatfish and the snail *Lunatia lewisii*, were rare in culture plots when gear was present. These organisms may actively avoid habitat complexity created by aquaculture gear. Holsman et al. (2006) found that subadult Dungeness crab (*Metacarcinus magister*, formerly *Cancer magister*) similarly avoid complex habitats, including on-bottom oyster culture, and preferentially use unstructured habitats during intertidal forays. For taxa adapted to unstructured habitat, complexity may hinder movement and reduce foraging efficiency (e.g., Holsman et al. 2010). The habitat value of unstructured areas to these taxa is substantial and should be considered along with any perceived positive habitat value of aquaculture gear to structure-oriented or crevice-dwelling fish and macroinvertebrates.

Effects of aquaculture on transient macrofauna did not persist after PVC tubes and nets were removed during grow-out. There was a significant difference between the culture plots for the last two aquaculture phases: gear present versus postgear ( $R = 0.160$ ,  $P = 0.003$ ), and the ANOSIM  $R$  value for this test



**Figure 9.** Two-dimensional nonmetric multidimensional scaling plot (NMDS) of scuba surveys at culture plots and reference areas when aquaculture gear was in place (gear present) and after gear was removed (postgear). The functional group vectors shown are those with  $P < 0.05$ . Stress = 18.08. Stress tested statistically significant with the Monte Carlo randomization approach ( $P = 0.03$ ). A goodness-of-fit Shepard plot showed good correlation between the ordination distances and the Bray–Curtis dissimilarities (linear fit  $R^2 = 0.877$ ). Vector loadings are shown for significant functional groups ( $P < 0.05$ ).

was the highest of all tests conducted, suggesting moderate ecological significance, which is corroborated by the NMDS plot in Figure 8. Moreover, when PVC tubes and nets were removed, the transient macrofauna community was no different from the pregear condition (ANOSIM  $R = -0.085$ ,  $P = 0.842$ ). These data suggest transient macrofauna communities associated with these intertidal beaches begin to recover to preaquaculture conditions within a few months of removal of the PVC tubes and nets.

Transient macrofaunal communities in reference areas were also significantly different between the gear-present and post-gear phases. The similar pattern observed in both culture plots and reference areas may be attributed at least in part to annual variation in species abundance and composition. Spatial and temporal variability can strongly influence transient macrofauna communities on a variety of scales (Jackson & Jones 1999, Hurst et al. 2004), and these changes can produce effects across trophic levels (Reum & Essington 2008). Reference areas in the current study may also be somewhat affected by removal of aquaculture structures between the gear-present and postgear phases through spillover effects (e.g., Ries & Sisk 2004). Culture plots and reference areas were 75–150 m apart. Previous work has demonstrated spillover effects on transient macrofauna from both natural (Almany 2004) and artificial structures (Helvey 2002).

Geoduck aquaculture practices did not affect diversity of macrofauna. No consistent differences in diversity of resident macrofauna were observed in the current study. Average diversity of transient macrofauna at culture plots when gear was present was slightly greater than at reference areas (but not significant), and diversity measures for the pregear and

postgear data were almost identical between habitat types. It is important to note that the Shannon index is based on relative instead of absolute abundance. This distinction is a potential limitation for a study such as the current one, which focuses on distinguishing between the raw abundance of species groups in different areas. Nevertheless, the results clearly contrast with previous work linking aquaculture disturbance with changes in diversity (Erbland and Ozbay [2008]; see review by Dumbauld et al. [2009]). Brown and Thuesen (2011) observed greater diversity of transient macrofauna associated with geoduck aquaculture gear in trapping surveys. However, taxa richness was low in that study, and results were driven by a large number of graceful crab—*Cancer (Metacarcinus) gracilis*—captured in the reference area. Overall, more organisms were captured in traps set in the reference area than in the geoduck aquaculture plots (Brown & Thuesen 2011).

Managers and stakeholders have raised concerns about potential effects of geoduck aquaculture practices on forage fish spawning habitat, particularly Pacific sand lance (*Ammodytes hexapterus*), which spawn on littoral beaches at high tidal levels (November to April [Penttila 2007]). Despite the presence of adult fish in excavation samples (Rogers site, October 2010), no evidence of spawning (i.e., eggs) was observed. It is possible that adult sand lance do not form winter aggregations in the same littoral habitats where spawning occurs (Quinn 1999). Moulton and Penttila (2000) suggest that spawning typically occurs at 2–2.75 m above MLLW, which is well above geoduck aquaculture operations and sampling in the current study (Table 1). No other adult forage fish, such as surf smelt (*Hypomesus pretiosus*) and herring (*Clupea pallasii*), or evidence of spawning activities were observed during the study. Although these results suggest negligible effects, the opportunistic sampling may be inadequate given spatiotemporal variability in spawning behavior, and additional targeted investigation is warranted to elucidate potential broader impacts on forage fish populations.

The current study provides insight into the response of resident and transient macrofauna to geoduck aquaculture practices. Taken together, these results indicate that changes in habitat complexity associated with geoduck aquaculture produce short-term effects (1–2 y) on intertidal beaches. However, it should be noted that the current study focused exclusively on diversity and abundance of fish and macroinvertebrate communities. Additional impacts might be demonstrated by considering different metrics, including growth. For example, Tallman and Forrester (2007) found that scup were 40% smaller in oyster cages relative to natural rocky areas, despite greater abundance of the species at aquaculture sites. Also, the current study focused on three isolated aquaculture operations over a single culture

cycle. Thus, it is not possible to extrapolate results to consider the cumulative effects of multiple culture cycles in a single location through repeated disturbance, or the landscape effects of a mosaic of adjacent aquaculture areas interspersed with other habitat types (see Dumbauld et al. 2009). Additional monitoring efforts and spatially explicit modeling work are required to develop an understanding of these phenomena, which are critical if this method of aquaculture continues to expand in the region. Moreover, the sampling used in the current study was not adequate to assess rare or patchy species, particularly salmonids. Scuba surveys and shoreline transects provide only a cursory appraisal of salmonid habitat use in this context, and given the contentious nature of salmon management in the region, rigorous assessment is critical. It is recommended that alternative sampling methods, such as beach seining, be used to evaluate use of geoduck aquaculture by out-migrating smolts.

Future research should focus on the issues just described, as well as on ecosystem effects on higher trophic levels. Nevertheless, the results of this study provide valuable insight into the ecological effects of geoduck aquaculture practices and add to a growing body of work describing the effects of anthropogenic disturbance on nearshore marine ecosystems. Most important, these data will aid regulatory authorities and resource managers in placing aquaculture-related disturbance in an appropriate context for decision making to balance the needs of stakeholders and environmental protection.

#### ACKNOWLEDGMENTS

The authors thank J. Eggers, N. Grose, M. Langness, F. Stevick, D. Todd, J. Toft, and others who assisted with scuba and collected field data in South Puget Sound. J. Olden consulted on aspects of the analyses. Helpful comments and assistance from K. Holsman and T. Essington greatly improved the manuscript. Valuable administrative assistance was provided by the Washington Sea Grant Program. This research was funded in part by the Washington state legislature, Washington Department of Ecology, Washington Department of Natural Resources, and Northwest Indian Fisheries Commission through the Shellfish Management Department of the Point No Point Treaty Council. Additional support to A.W.E.G. was provided by a Castagna Student Grant for Applied Research (2009; National Shellfisheries Association), and to G.R.V. by the U.S. Geological Survey (USGS). The views expressed herein are those of the authors and do not necessarily reflect the views of the funding agencies, excepting USGS. Any use of trade, firm, or product name is for descriptive purposes only and does not imply endorsement by the U.S. government.

#### LITERATURE CITED

- Almany, G. R. 2004. Differential effects of habitat complexity, predators and competitors on abundance of juvenile and adult coral reef fishes. *Oecologia* 141:105–113.
- Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* 26:32–46.
- Anderson, M. J. 2006. Distance-based tests for the homogeneity of multivariate dispersions. *Biometrics* 62:245–253.
- Bates, D. M. & M. Maechler. 2010. lme4. Linear mixed-effects models using S4 classes. Available at: <http://lme4.r-forge.r-project.org/>.
- Bernard, F. R. 1983. Catalogue of the living Bivalvia of the eastern Pacific Ocean: Bering Strait to Cape Horn. Ottawa, Ontario: Department of Fisheries and Oceans. 102 pp.
- Bradbury, A., B. Sizemore, D. Rothaus & M. Ulrich. 2000. Stock assessment of subtidal geoduck clams (*Panopea abrupta*) in Washington. Olympia, WA: Marine Resources Unit, Washington Department of Fish and Wildlife 57 pp.
- Bray, J. R. & J. T. Curtis. 1957. An ordination of the upland forest communities of southern Wisconsin. *Ecol. Monogr.* 48:35–49.

- Brown, R. A. & E. V. Thuesen. 2011. Biodiversity of mobile benthic fauna in geoduck (*Panopea generosa*) aquaculture beds in southern Puget Sound, Washington. *J. Shellfish Res.* 30:771–776.
- Burns, R. E. 1985. The shape and form of Puget Sound. Puget Sound Books, Washington Sea Grant Program, University of Washington. Seattle, WA: University of Washington Press. 100 pp.
- Caswell, H. & J. E. Cohen. 1991. Disturbance, interspecific interaction and diversity in metapopulations. *Biol. J. Linn. Soc. Lond.* 42:193–218.
- Clarke, K. R. 1993. Non-parametric multivariate analyses of changes in community structure. *Aust. J. Ecol.* 18:117–143.
- Clarke, K. R. & R. N. Gorley. 2001. PRIMERv5: user manual/tutorial. Plymouth, UK: PRIMER-E. 91 pp.
- Connell, J. H. 1978. Diversity in tropical rain forests and coral reefs. *Science* 199:1302–1309.
- Costa-Pierce, B. 2002. Ecology as the paradigm for the future of aquaculture. In: B. Costa-Pierce, editor. Ecological aquaculture: the evolution of the blue revolution. Oxford, UK: Blackwell Science. pp. 339–372.
- Crowder L. B. & W. E. Cooper. 1982. Habitat structural complexity and the interaction between bluegills and their prey. *Ecology* 63:1802–1813.
- Davis, N., G. R. VanBlaricom & P. K. Dayton. 1982. Man-made structures on marine sediments: effects on adjacent benthic communities. *Mar. Biol.* 70:295–303.
- Dealteris, J. T., B. D. Kilpatrick & R. B. Rheault. 2004. A comparative evaluation of the habitat value of shellfish aquaculture gear, submerged aquatic vegetation and a non-vegetated seabed. *J. Shellfish Res.* 23:867–874.
- Dethier, M. N. 2005. Spatial patterns and temporal trends in shoreline biota in Puget Sound: analyses of data collected through 2004. Nearshore Habitat Program. Olympia, WA: Washington Department of Natural Resources. 27 pp.
- Dubois, S., J. C. Marin-Leal, M. Ropert & S. Lefebvre. 2007. Effects of oyster farming on macrofaunal assemblages associated with *Lanice conchilega* tubeworm populations: a trophic analysis using natural stable isotopes. *Aquaculture* 271:336–349.
- Dumbauld, B. R., J. L. Ruesink & S. S. Rumrill. 2009. The ecological role of bivalve shellfish aquaculture in the estuarine environment: a review with application to oyster and clam culture in West Coast (USA) estuaries. *Aquaculture* 290:196–223.
- Erbland, P. J. & G. Ozbay. 2008. Comparison of the macrofaunal communities inhabiting a *Crassostrea virginica* oyster reef and oyster aquaculture gear in Indian River Bay, Delaware. *J. Shellfish Res.* 27:757–768.
- Ferraro, S. P. & F. A. Cole. 1990. Taxonomic level and sample size sufficient for assessing pollution impacts on the Southern California Bight macrobenthos. *Mar. Ecol. Prog. Ser.* 67:251–262.
- FAO. 2012. Global aquaculture production [online query]. Available at: <http://www.fao.org/fishery/aquaculture/en>. Accessed June 19, 2013.
- Genkai-Kato, M. 2007. Macrophyte refuges, prey behavior and trophic interactions: consequences for lake water clarity. *Ecol. Lett.* 10:105–114.
- Glasby, T. M. & A. J. Underwood. 1996. Sampling to differentiate between pulse and press perturbations. *Environ. Monit. Assess.* 42:241–252.
- Goodwin, C. & B. Pease. 1987. The distribution of geoduck (*Panopea abrupta*) size, density, and quality in relation to habitat characteristics such as geographic area, water depth, sediment type, and associated flora and fauna in Puget Sound, Washington. Olympia, WA: Washington Department of Fisheries, Shellfish Division. 44 pp.
- Grabowski, J. H. 2004. Habitat complexity disrupts predator–prey interactions but not the trophic cascade on oyster reefs. *Ecology* 85:995–1004.
- Grabowski, J. H. & S. P. Powers. 2004. Habitat complexity mitigates trophic transfer on oyster reefs. *Mar. Ecol. Prog. Ser.* 277:291–295.
- Healey, M. C. 1979. Detritus and juvenile salmon production in the Nanaimo estuary. I: production and feeding rates of juvenile chum salmon (*Oncorhynchus keta*). *J. Fish. Res. Board Can.* 36:488–496.
- Helvey, M. 2002. Are southern California oil and gas platforms essential fish habitat? *ICES J. Mar. Sci.* 59:S266–S271.
- Holsman, K. K., P. S. McDonald & D. A. Armstrong. 2006. Intertidal migration and habitat use by subadult Dungeness crab *Cancer magister* in a NE Pacific estuary. *Mar. Ecol. Prog. Ser.* 308:183–195.
- Holsman, K. K., P. S. McDonald, P. A. Barretero & D. A. Armstrong. 2010. Restoration through eradication? Removal of an invasive bioengineer restores some habitat function for a native predator. *Ecol. Appl.* 20:2249–2262.
- Hurst, T. P., K. A. McKown & D. O. Conover. 2004. Interannual and long-term variation in the nearshore fish community of the mesohaline Hudson River estuary. *Estuaries* 27:659–669.
- Inglis, G. J. & N. Gust. 2003. Potential indirect effects of shellfish culture on the reproductive success of benthic predators. *J. Appl. Ecol.* 40:1077–1089.
- Jackson, G. & G. K. Jones. 1999. Spatial and temporal variation in nearshore fish and macroinvertebrate assemblages from a temperate Australian estuary over a decade. *Mar. Ecol. Prog. Ser.* 182:253–268.
- Kruskal, J. B. & M. Wish. 1978. Multidimensional scaling. Beverly Hills, CA: Sage. 93 pp.
- Lapointe, L. & E. Bourget. 1999. Influence of substratum heterogeneity scales and complexity on a temperate epibenthic marine community. *Mar. Ecol. Prog. Ser.* 189:159–170.
- Lenihan, H. S. 1999. Physical–biological coupling on oyster reefs: how habitat structure influences individual performance. *Ecol. Monogr.* 69:251–275.
- Mackey, R. L. & D. J. Currie. 2001. The diversity–disturbance relationship: is it generally strong and peaked? *Ecology* 82:3479–3492.
- Mattsson, J. & O. Linden. 1983. Benthic macrofauna succession under mussels, *Mytilus edulis* L. (Bivalvia), cultured on hanging long-lines. *Sarsia* 68:97–102.
- McCullagh, P. & J. A. Nelder. 1989. Generalized linear models, 2<sup>nd</sup> edition. London: Chapman and Hall. 536 pp.
- McCune, B., J. B. Grace & D. L. Urban. 2002. Analysis of ecological communities. Gleneden Beach, OR: MjM Software Design. 300 pp.
- McGarigal, K., S. A. Cushman & S. G. Stafford. 2000. Multivariate statistics for wildlife and ecology research. New York: Springer. 283 pp.
- Moulton, L. & D. E. Penttila. 2000. San Juan County forage fish assessment project: forage fish spawning distribution in San Juan County and protocols for sampling intertidal and nearshore regions final report. Mount Vernon, WA: San Juan County Marine Resource Committee and Northwest Straits Commission. 36 pp.
- Penttila, D. 2007. Marine forage fishes in Puget Sound. Puget Sound Nearshore Partnership report no. 2007-03. Seattle, WA: Seattle District, U.S. Army Corps of Engineers. 23 pp.
- Pickett, S. T. A. & P. S. White, editors. 1985. The ecology of natural disturbance and patch dynamics. New York: Academic Press. 472 pp.
- Powers, M. J., C. H. Peterson, H. C. Summerson & S. P. Powers. 2007. Macroalgal growth on bivalve aquaculture netting enhances nursery habitat for mobile invertebrates and juvenile fishes. *Mar. Ecol. Prog. Ser.* 339:109–122.
- Quinn, T. 1999. Habitat characteristics of an intertidal aggregation of Pacific sand lance (*Ammodytes hexapterus*) at a North Puget Sound beach in Washington. *Northwest Sci.* 73:44–49.
- Quintino, V., A. Azevedo, L. Magalhaes, L. Sampaio, R. Freitas, A. M. Rodrigues & M. Elliott. 2012. Indices, multispecies and synthesis descriptors in benthic assessments: intertidal organic enrichment from oyster farming. *Estuar. Coast. Shelf Sci.* 110:190–201.
- R Development Core Team. 2011. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Available at: <http://www.R-project.org>.
- Reum, J. C. P. & T. E. Essington. 2008. Seasonal variation in guild structure of the Puget Sound demersal fish community. *Estuaries Coasts* 31:790–801.

- Ries, L. & T. D. Sisk. 2004. A predictive model of edge effects. *Ecology* 85:2917–2962.
- Shannon, C. E. 1948. A mathematical theory of communication. *Bell Syst. Tech. J.* 27:379–423, 623–656.
- Shannon, C. E. & W. Weaver. 1949. The mathematical theory of communication. Champaign, IL: University of Illinois Press. 117 pp.
- Simenstad, C. A. & K. L. Fresh. 1995. Influence of intertidal aquaculture on benthic communities in Pacific Northwest estuaries: scales of disturbance. *Estuaries* 18:43–70.
- Simenstad, C. A., K. L. Fresh & E. O. Salo. 1982. The role of Puget Sound and Washington coastal estuaries in the life history of Pacific salmon: an unappreciated function. In: V. S. Kennedy, editor. *Estuarine comparisons*. New York: Academic Press. pp. 343–364.
- Simenstad, C. A., C. T. Tanner, R. M. Thom & L. L. Conquest. 1991. *Estuarine habitat assessment protocol*. Seattle, WA: U.S. Environmental Protection Agency, Region 10. 201 pp.
- Sobocinski, K. L., J. R. Cordell & C. A. Simenstad. 2010. Effects of shoreline modifications on supratidal macroinvertebrate fauna on Puget Sound, Washington beaches. *Estuaries Coasts* 33:699–711.
- Spencer, B. E., M. J. Kaiser & D. B. Edwards. 1997. Ecological effects of intertidal Manila clam cultivation: observations at the end of the cultivation phase. *J. Appl. Ecol.* 34:444–452.
- Tallman, J. C. & G. E. Forrester. 2007. Oyster grow-out cages function as artificial reefs for temperate fishes. *Trans. Am. Fish. Soc.* 136:790–799.
- VanBlaricom, G. R., J. L. Eccles, J. D. Olden & P. S. McDonald. 2015. Ecological effects of the harvest phase of geoduck (*Panopea generosa* Gould, 1850) aquaculture on infaunal communities in southern Puget Sound, Washington. *J. Shellfish Res.* 34:171–187.
- Warwick, R. M. & K. R. Clarke. 1993. Increased variability as a symptom of stress in marine communities. *J. Exp. Mar. Biol. Ecol.* 172:215–226.
- Wolfson, A., G. R. VanBlaricom, N. Davis & G. Lewbel. 1979. The marine life of an offshore oil platform. *Mar. Ecol. Prog. Ser.* 1:81–89.
- Woodin, S. A. 1981. Disturbance and community structure in a shallow water sand flat. *Ecology* 62:1052–1066.
- Young, C. D. 2009. Shoaling behavior as a tool to understand microhabitat use by juvenile chum salmon, *Oncorhynchus keta*. MS thesis, School of Aquatic and Fishery Sciences, University of Washington. 83 pp.
- Zar, J. H. 2010. *Biostatistical analysis*. Upper Saddle River, NJ: Prentice-Hall/Pearson. 960 pp.